

Spontaneous Symmetry Breaking in Directed Percolation with Many Colors: Differentiation of Species in the Gribov Process

H.K. JANSSEN

*Institut für Theoretische Physik III,
Heinrich-Heine-Universität, 40225 Düsseldorf, Germany
(February 1, 2008)*

A general field theoretic model of directed percolation with many colors that is equivalent to a population model (Gribov process) with many species near their extinction thresholds is presented. It is shown that the multicritical behavior is always described by the well known exponents of Reggeon field theory. In addition this universal model shows an instability that leads in general to a total asymmetry between each pair of species of a cooperative society.

PACS numbers: 64.60.Ak, 05.40.+j, 64.60.Ht

It is well known that the Gribov process [1,2] (Reggeon field theory [3–5], stochastic Schlögl model [6]) is a stochastic multiparticle process which describes the essential features of a vast number of growth phenomena of populations near their extinction threshold and without exploitation of the environment. The transition between survival and extinction of the population is a nonequilibrium continuous phase transition phenomenon and is characterized by universal scaling laws. The Gribov process belongs to the universality class of local growth processes with absorbing states [7,8] such as directed percolation [9–11], the contact process [12–14], and certain cellular automata [15,16], and is relevant to a vast range of models in physics, chemistry, biology, and sociology. I have been studying this transition for some time by renormalized field theoretic methods [7]. In this letter I will show that the Gribov process with many interacting species (directed percolation with many colors) also belongs to this universality class. In addition this multicolored directed percolation exhibits a new interesting phenomenon: even if the dynamics of the different colors is modeled symmetrically, there exists an instability which leads to a differentiation of the species in their active state. This color symmetry breaking leads to many different stable forms of a cooperative society.

The multiparticle processes under consideration are characterized by the following four principles:

1. Self-reproduction (“birth”) is error free, but may be different for each species; particles can be annihilated with rates depending on the species (“death”).
2. The particles interact with species-dependent couplings (“competition” and “cooperation”).
3. Particles diffuse in a d -dimensional space (“motion”).
4. The states where some species are extinct are absorbing.

In the language of chemistry, the system corresponds to an autocatalytic reaction scheme of the form $X_\alpha \leftrightarrow 2X_\alpha$, $X_\alpha \rightarrow 0$, $X_\alpha + X_\beta \rightarrow kX_\alpha + lX_\beta$, where the last reaction subsumes the interactions of species with different colors α, β , and $k, l = 0, 1$. A description in terms of particle densities typically arises from a coarse-graining procedure where a large number of microscopic degrees of freedom are averaged out. The influence of these is simply modeled as Gaussian noise-terms in a Langevin equation which have to respect the absorbing state condition. Stochastic reaction-diffusion equations for the particle densities $n(\mathbf{r}, t) = (n_1(\mathbf{r}, t), n_2(\mathbf{r}, t), \dots)$ in accordance with the four principles given above are constructed as

$$\partial_t n_\alpha(\mathbf{r}, t) = \lambda_\alpha \nabla^2 n_\alpha(\mathbf{r}, t) + R_\alpha(n(\mathbf{r}, t)) n_\alpha(\mathbf{r}, t) + \zeta_\alpha(\mathbf{r}, t), \quad (1)$$

where the first term on the right hand side models the (diffusive) motion, and the R_α are the reproduction rates of the particles with color α . These deterministic terms are constructed proportional to n_α in order to ensure the existence of an absorbing state for each species. Expanding the rates R_α in powers of n results in

$$R_\alpha(n(\mathbf{r}, t)) = -\lambda_\alpha \left[\tau_\alpha + \frac{1}{2} \sum_\beta g_{\alpha\beta} n_\beta(\mathbf{r}, t) \right] \quad (2)$$

up to subleading terms. The Gaussian noises $\zeta_\alpha(\mathbf{r}, t)$ must also respect the absorbing state condition, whence

$$\langle \zeta_\alpha(\mathbf{r}, t) \zeta_\beta(\mathbf{r}', t') \rangle = \lambda_\alpha g_{\alpha\beta} \delta_{\alpha\beta} n_\alpha(\mathbf{r}, t) \times \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \quad (3)$$

also up to subleading contributions. In this letter I only consider color-independent diffusion constants $\lambda_\alpha = \lambda$. Then, for stability, it must be demanded $\sum_{\alpha,\beta} n_\alpha g_{\alpha\beta} n_\beta \geq 0$ for all $n_\alpha \geq 0$. The “temperature” variables τ_α measure the difference of the rates of death and birth of the species α . Thus the temperatures may be positive or negative. I am interested in the case where all

$\tau_\alpha \approx 0$ (up to fluctuation corrections) defining the multicritical region. Under these conditions all the species live on the border of extinction.

The equations of motion (1) show (absorbing) homogeneous steady state solutions with any combination of extinct species. In the infinite volume limit one finds transitions to steady states with $n_\alpha \geq 0$ which are given in mean field theory as the solutions of $\sum_\beta g_{\alpha\beta} n_\beta = -2\tau_\alpha$ if $n_\alpha > 0$. This set of equations in general offers multiple different solutions if some or all of the τ_α are negative. Each of these solutions defines a different homogeneous phase of the active system. In some of these phases only one of the species is alive, in the other phases several species act cooperatively. In the space spanned by the temperature variables τ_α , the phases are separated by surfaces of first or second order transitions where a species becomes extinct. All the critical surfaces meet together in a multicritical point where all temperatures are zero. Take e.g. the simplest case of a totally symmetric three-species system with $g_{\alpha\beta} = g$ and $\tau_\alpha < 0$ for $\alpha, \beta = 1, 2, 3$. The ternary phase diagram is shown in Fig.1. In this case there only exist phases where one species is nonzero and the two others are suppressed. Here stable cooperative phases with more than one species are not possible.

In general, this mean field picture continuously depends on the coupling constants $g_{\alpha\beta}$. In the following I will show that in spatial dimensions lower than 4, the stochastic equations (1-3) produce universal macroscopic steady state equations which correspond to only a discrete set of choices for the coupling matrix $\mathbf{g}$ that I call strategies. The stable ones are distinguished by a total asymmetry between the members of each pair of species, in a mean field picture: $g_{\alpha\beta}$ or $g_{\beta\alpha}$ is zero. This implies that only one of these partners directly influences the evolution of the other. These stable strategies with differentiated species permit cooperative phases with more than one species nonzero and continuous phase transitions. All the continuous phase transitions of each strategy are characterized by the directed percolation critical exponents of the one-species Gribov process.

In the following I will be interested in the behavior of correlation- and response functions in the vicinity of the multicritical point where all τ_α are small. The critical properties are extracted using renormalized field theory [17–22] in conjunction with an ε -expansion about the upper critical dimension $d_c = 4$. It is then convenient to recast the Langevin equation as a dynamical functional [23,24]

$$\mathcal{J}[\tilde{s}, s] = \int dt d^d r \sum_\alpha \left\{ \tilde{s}_\alpha \left[\partial_t + \lambda_\alpha (\tau_\alpha - \nabla^2) + \frac{1}{2} \sum_\beta g_{\alpha\beta} s_\beta - \frac{1}{2} g_{\alpha\alpha} \tilde{s}_\alpha \right] s_\alpha \right\}, \quad (4)$$

where $s_\alpha(\mathbf{r}, t) \sim n_\alpha(\mathbf{r}, t)$ are rescaled density variables which ensure that $g_{\alpha\alpha} = g_\alpha$. The $\tilde{s}_\alpha(\mathbf{r}, t)$ denote the response fields (the MSR auxiliary variables [25]). Correlation- and response functions can now be expressed as functional averages of monomials of s and $\tilde{s}$ with weight $\exp\{-\mathcal{J}\}$. The responses are defined with respect to additional local particle sources $\tilde{h}_\alpha(\mathbf{r}, t) \geq 0$ in the equation of motion (1). By these sources a further term $-\sum_\alpha \tilde{h}_\alpha \tilde{s}_\alpha$ is added to the integrand of the dynamic functional $\mathcal{J}$ (4). Coupling constants invariant under the rescaling $s_\alpha \rightarrow c_\alpha s_\alpha$, $\tilde{s}_\alpha \rightarrow c_\alpha^{-1} \tilde{s}_\alpha$ are given by $f_{\alpha\beta} = g_{\alpha\beta} g_\beta$. The scaling by the usual convenient length μ^{-1} and time scale $(\lambda\mu^2)^{-1}$, respectively, shows $\tilde{s}_\alpha \sim s_\alpha \sim \mu^{d/2}$, $f_{\alpha\beta} \sim \mu^\varepsilon$ where $\varepsilon = 4 - d$ which signals $d_c = 4$ to be the upper critical dimension. A further inspection of (4) and a glance at the perturbation expansion shows that the functional $\mathcal{J}$ includes all relevant couplings and is thus renormalizable.

To determine the renormalizations, one has to calculate the primitively divergent vertex functions $\Gamma_{\alpha,\alpha}$, $\Gamma_{\alpha\alpha,\alpha} = \Gamma_{\alpha,\alpha\alpha}$, $\Gamma_{\alpha,\alpha\beta}$ with nonnegative μ -dimensions. Here the first and the last part of indices denotes the amputated $\tilde{s}$ - and s -legs, respectively. Dimensional regularization and minimal renormalization is used. The ε -poles are absorbed by multiplicative renormalizations of the fields and couplings which I choose in the form

$$\begin{aligned} s_\alpha &\rightarrow \mathring{s}_\alpha = \sqrt{Z_s^{(\alpha)}} s_\alpha, & \tilde{s}_\alpha &\rightarrow \mathring{\tilde{s}}_\alpha = \sqrt{Z_s^{(\alpha)}} \tilde{s}_\alpha, \\ \lambda_\alpha &\rightarrow \mathring{\lambda}_\alpha = \frac{Z_\lambda^{(\alpha)}}{Z_s^{(\alpha)}} \lambda_\alpha, & \tau_\alpha &\rightarrow \mathring{\tau}_\alpha = \frac{Z_\tau^{(\alpha)}}{Z_\lambda^{(\alpha)}} \tau_\alpha, \\ f_{\alpha\beta} &\rightarrow \mathring{f}_{\alpha\beta} = G_\varepsilon^{-1} \mu^\varepsilon \frac{Z_u^{(\alpha\beta)}}{Z_s^{(\beta)} Z_\lambda^{(\alpha)} Z_\lambda^{(\beta)}} u_{\alpha\beta}. \end{aligned} \quad (5)$$

Here $G_\varepsilon = \Gamma(1 + \varepsilon/2) / (4\pi)^{d/2}$ is a convenient constant. From the topology of the diagrams of the perturbation expansion it is easily seen that the vertex functions do not depend on coupling constants with indices other than the ones of the vertex functions itself. Therefore the renormalization factors $Z_i^{(\alpha)}$ with $i = (s, \tau, \lambda)$ and $Z_u^{(\alpha)} := Z_u^{(\alpha\alpha)}$ that are determined from $\Gamma_{\alpha,\alpha}$ and $\Gamma_{\alpha\alpha,\alpha}$ are already known from the simple Gribov process. Thus the renormalization group flows of the fields and parameters with exception of the $g_{\alpha\beta}$ with $\alpha \neq \beta$ do not depend on the coupling between different colors. It follows at once that all the critical exponents are given by the usual directed percolation exponents and the fixed point values of the “diagonal” renormalized couplings $u_{\alpha\alpha} = u_\alpha$ are all independent from the color α .

The flow of the couplings between different colors follows from the $Z_u^{(\alpha\beta)}$ with $\alpha \neq \beta$. The symmetric fixed

point might be attractive. To disprove that I have calculated these renormalizations up to two loop order for the special case $\lambda_\alpha = \lambda$, $u_\alpha := u_{\alpha\alpha} = u$ with the result

$$\begin{aligned} Z_u^{(\alpha\beta)} = 1 + \frac{1}{4\varepsilon} (6u + u_{\alpha\beta} + u_{\beta\alpha}) + \frac{1}{16\varepsilon} \left[\left(\frac{36}{\varepsilon} - \frac{19}{2} \right) u^2 \right. \\ \left. + \left(\frac{8}{\varepsilon} - \frac{5}{4} \right) uu_{\beta\alpha} + \left(\frac{1}{\varepsilon} - \frac{3}{2} \ln \frac{4}{3} \right) (u_{\alpha\beta}^2 + u_{\beta\alpha}^2) \right. \\ \left. + \left(\frac{2}{\varepsilon} - 1 \right) u_{\alpha\beta} u_{\beta\alpha} + \left(\frac{8}{\varepsilon} - \frac{9}{4} + 3 \ln \frac{4}{3} \right) u_{\alpha\beta} u \right]. \end{aligned}$$

Collecting together these and the known results for the other renormalization factors, one easily gets the flow equation for the renormalized coupling constants

$$\mu \frac{\partial}{\partial \mu} u_{\alpha\beta} = \beta_{\alpha\beta} \quad (6)$$

with the Wilson-functions

$$\begin{aligned} \beta_{\alpha\beta} = \left[-\varepsilon + u + \frac{1}{4} (u_{\alpha\beta} + u_{\beta\alpha}) - \frac{1}{8} u_{\alpha\beta} u_{\beta\alpha} \right. \\ \left. - \left(\frac{97}{128} + \frac{53}{64} \ln \frac{4}{3} \right) u^2 - \left(\frac{9}{32} - \frac{3}{8} \ln \frac{4}{3} \right) uu_{\alpha\beta} \right. \\ \left. - \frac{5}{32} uu_{\beta\alpha} - \frac{3}{16} \ln \frac{4}{3} (u_{\alpha\beta}^2 + u_{\beta\alpha}^2) + \dots \right] u_{\alpha\beta}. \quad (7) \end{aligned}$$

For $\alpha = \beta$, the equation $\beta_{\alpha\alpha} =: \beta(u_\alpha) = 0$ yields the nontrivial stable fixed point value $u_* = \frac{2}{3}\varepsilon \{1 + (\frac{169}{288} + \frac{53}{144} \ln \frac{4}{3})\varepsilon + \dots\}$. For each pair of species with colors $\alpha \neq \beta$, fixed point solutions of (6) with at least one zero coupling are either given by $u_{\alpha\beta*} = u_{\beta\alpha*} = 0$, or by the pair of solutions $u_{\alpha\beta*} = (1 + \varepsilon_{\alpha\beta}) w_*$ with $\varepsilon_{\alpha\beta} = -\varepsilon_{\beta\alpha} = \pm 1$ and $w_* = u_* + O(\varepsilon^3)$. To analyze the case with $u_{\alpha\beta*}$ and $u_{\beta\alpha*}$ both nonzero, one sets $u_{\alpha\beta*} = u_* + v_{\alpha\beta}$. To first order in ε one finds a fixed point line $v_{\alpha\beta} + v_{\beta\alpha} = 0$. But using the higher order terms of (7), one gets $v_{\alpha\beta} = 0 + O(\varepsilon^2)$. Then it can be easily seen that $v_{\alpha\beta}$ must be zero order for order in the ε -expansion. This proves that in this case the fixed point is indeed symmetric $u_{\alpha\beta*} = u_{\beta\alpha*} = u_*$. Thus I have found from this analysis that for each pair of species independent of the other ones three types of fixed points for the couplings exist: both couplings vanish (decoupled), or only one of the couplings is nonzero (asymmetric), or both couplings are nonzero and equal to the diagonal one (symmetric).

The stability of the various fixed points is analyzed by the calculation of the eigenvalues of the matrix of derivatives of the Wilson-functions $\beta_{\alpha\beta}$, $\beta_{\beta\alpha}$. This analysis yields as stable fixed points only the asymmetric ones. The decoupled fixed points are doubly unstable, the two instability exponents are equal to $\omega_d = \frac{1}{2}u_* (1 - \frac{9}{8}u_*) + \dots = \frac{\varepsilon}{3} (1 - \varepsilon 0.0573 + \dots)$. The symmetric fixed points have one instability exponent $\omega_s = (1 - 3 \ln \frac{4}{3}) u_*^2 + \dots = \varepsilon^2 0.0609 + \dots$. I especially note

the very slow crossover from the symmetric to the asymmetric fixed point.

To get a simple qualitative impression of the behavior of the dynamic system at its asymmetric fixed points, I consider, in the active region with all $-\tau_\alpha =: g r_\alpha \geq 0$, a renormalized mean-field theory of the equations (1,2) for spatially homogeneous densities $\bar{n}_a$ and set $g_{\alpha\alpha} = g$, $g_{\alpha\beta} = g(1 + \varepsilon_{\alpha\beta})$ and $\sum_\alpha \bar{n}_a = \bar{n}$:

$$\partial_t \bar{n}_a = \lambda g \left(r_\alpha - \frac{1}{2} \bar{n} - \frac{1}{2} \sum_\beta \varepsilon_{\alpha\beta} \bar{n}_\beta \right) \bar{n}_\alpha. \quad (8)$$

The decision which of all the $\varepsilon_{\alpha\beta} = -\varepsilon_{\beta\alpha}$ are ± 1 defines a stable strategy. The strategies may be pictured by a strategy-diagram. Each corner of such a diagram represents a color α . Consider now each pair (α, β) of corners, they are connected by a line with an arrow pointing at α if $\varepsilon_{\alpha\beta} = 1$ or vice versa if $\varepsilon_{\alpha\beta} = -1$. The arrows therefore represent the “pressure” on the reproduction rate resulting from one species to another. The diagrams for the stable strategies are therefore fully connected by such directed lines. For a system with three species e.g. two stable strategies are possible, a cyclic and a hierarchic one (upper part of Fig.2). In (8) the influence on the reproduction rates of the society as a whole is described by the common “pressure” $p = g\bar{n}/2$. If now $\varepsilon_{\alpha\beta} = 1$ e.g., the part of p resulting from the species β on the reproduction of α is doubled whereas the part of the common pressure resulting from the species α on the reproduction of β is fully compensated. As a result there exist many cooperative phases with more than one species as stationary solutions of the mean field equations of motion. As an example the phase diagrams for the stable strategies of a system of three species is shown in Fig.2 (lower part).

Now the unstable strategies can be pictured as diagrams with disconnected corners for a pair (α, β) in the uncoupled case $g_{\alpha\beta} = 0$ (then there is no pressure from α on β and vice versa), and connected corners but without direction for a pair (α, β) in the symmetric case $g_{\alpha\beta} = g_{\beta\alpha}$ (then the species α, β do influence their reproduction rates only by means of the common pressure p). It is easily seen that in the fully symmetric case (for three species see Fig.1) more than one species can coexist only if their temperatures $\tau_\alpha = -g r_\alpha$ are all equal. If for one color α the parameter r_α is slightly increased, the species α suppresses all the others, if r_α is decreased, it becomes extinct by the common pressure of all the others.

Summarily I have shown that also in stochastic multispecies models of populations that develops near the extinction threshold of all the species and therefore have many absorbing states, the critical properties at the multicritical point and at all continuous transitions are governed by the well known Gribov process (Reggeon field theory) exponents (for a previous simulational result on

a two-species system which seems to agree with my findings see [26]). In other regions of the phase diagram of course more complicated critical behavior may arise such as multicritical points with different scaling exponents (see the recent work of Bassler and Brown [27]). The models considered here have many absorbing states (each combination of species may be extincted irrespectively of the other ones) and therefore are different from models which were considered by Grinstein et al. [28] where it was shown that multispecies systems with one absorbing state belong to the Gribov universality class. In addition I have shown that the universal properties of interspecies correlations and the phase diagram are determined by totally asymmetric fixed point values of the renormalized interspecies coupling constants. This eventually leads to a system working cooperatively. It is interesting that the asymmetry between the pairs of species seems to be the condition for this cooperation near extinction. The model considered here is a simple but universal model of such a cooperative society and should therefore have many applications in all fields of natural science up to sociology.

ACKNOWLEDGMENTS

I thank S. Theiss for a critical reading of the manuscript. This work has been supported in part by the SFB 237 of the Deutsche Forschungsgemeinschaft.

-
- [1] P. Grassberger and K. Sundermeyer, Phys. Lett. **77B**, 220 (1978).
 - [2] P. Grassberger and A. de La Torre, Ann. Phys. (New York), 373 (1979).
 - [3] V. N. Gribov, Sov. Phys. JETP **26**, 414 (1968).
 - [4] V. N. Gribov and A. A. Migdal, Sov. Phys. JETP **28**, 784 (1969).
 - [5] M. Moshe, Phys. Rep. C **37**, 255 (1978).
 - [6] F. Schlögl, Z. Phys. **225**, 147 (1972).
 - [7] H. K. Janssen, Z. Phys. B **42**, 151 (1981).
 - [8] P. Grassberger, Z. Phys. B **47**, 365 (1982).
 - [9] S. R. Broadbent and J. M. Hammersley, Proc. Camb. Philos. Soc. **53**, 629 (1957).

- [10] J. L. Cardy and R. L. Sugar, J. Phys. A: Math. Gen. **13**, L423 (1980).
- [11] S. P. Obukhov, Physica A **101**, 145 (1980).
- [12] T. E. Harris, Ann. Prob. **2**, 969 (1974).
- [13] T. M. Liggett, *Interacting Particle Systems*, (Springer, New York, 1985).
- [14] I. Jensen and R. Dickman, Physica A **203**, 175 (1994).
- [15] W. Kinzel, in *Percolation Structures and Processes*, edited by G. Deutsch, R. Zallen, and J. Adler, (Hilger, Bristol, 1983).
- [16] W. Kinzel, Z. Phys. B **58**, 229 (1985).
- [17] D. J. Amit, *Field Theory, the Renormalization Group and Critical Phenomena*, (World Scientific, Singapore, 1984).
- [18] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, 2nd revised edition, (Clarendon, Oxford, 1993).
- [19] R. Bausch, H. K. Janssen, and H. Wagner, Z. Phys. B **24**, 113 (1976).
- [20] C. DeDominicis and L. Peliti, Phys. Rev. B **18**, 353 (1978).
- [21] H. K. Janssen, in *Dynamical Critical Phenomena and Related Topics (Lecture Notes in Physics, Vol. 104)*, edited by C. P. Enz, (Springer, Heidelberg, 1979).
- [22] H. K. Janssen, in *From Phase Transitions to Chaos*, edited by G. Györgyi, I. Kondor, L. Sasvári, and T. Tél, (World Scientific, Singapore, 1992).
- [23] C. DeDominicis, J. Phys. (France) Colloq. **37**, C247 (1976).
- [24] H. K. Janssen, Z. Phys. B **23**, 377 (1976).
- [25] P. C. Martin, E. D. Siggia, and H. H. Rose, Phys. Rev. A **8**, 423 (1973).
- [26] S. Cornell, M. Droz, R. Dickman, and M. C. Marques, J. Phys. A: Math. Gen. **24**, 5605 (1991).
- [27] K. E. Bassler and D. A. Brown, Phys. Rev. Lett. **77**, 4094 (1996).
- [28] G. Grinstein, Z-W. Lai and D. A. Browne, Phys. Rev. A **40**, 4820 (1989).

FIG. 1. Symmetric strategy of a three-species-system and its ternary phase diagram in the plane $\tau_1 + \tau_2 + \tau_3 = -r < 0$. The numbers α at the corners denote the points where the corresponding $\tau_\alpha = -r$ and the other temperatures are zero. Dashed lines indicate discontinuous transitions between the one-species phases.

FIG. 2. Cyclic and hierarchic strategy of a three-species system and their ternary phase diagrams. The numbers inside the phase diagrams denote the colors of the living species. Solid lines indicate continuous transitions.



